

Front

You should immediately contact your doctor or healthcare provider if you think you are having a hypersensitivity reaction to Abacavir/Lamivudine tablets. Write your doctor's details below:

Doctor:

Tel:

If your doctor or healthcare provider is not available, you must urgently seek alternative medical advice (e.g. the emergency unit of the nearest hospital).

For general Abacavir/Lamivudine information enquiries, contact Generics (UK) Ltd T/A Mylan Tel: + 44 1707 853000.

IMPORTANT - ALERT CARD

Abacavir/Lamivudine tablets

Carry this card with you at all times

Back

Patients taking Abacavir may develop a hypersensitivity reaction (serious allergic reaction) which can be life threatening if treatment with Abacavir is continued.

CALL YOUR DOCTOR OR HEALTH CARE PROVIDER IMMEDIATELY for advice on whether you should stop taking Abacavir/Lamivudine if:

- 1) you get a skin rash **OR**
- 2) you get one or more symptoms from at least **TWO** of the following groups

Front

- fever
- shortness of breath, sore throat or cough
- nausea or vomiting or diarrhoea or abdominal pain
- severe tiredness or achiness or generally feeling ill

If you have discontinued Abacavir/Lamivudine due to this reaction, **YOU MUST NEVER TAKE** Abacavir/Lamivudine, or any medicine containing abacavir again as **within hours** you may experience a life-threatening lowering of your blood pressure or death.

(see reverse of card)

 Mylan

Page 2

Page 3