

Emergency contact information:

Emergency Prescriber Contact:

Telephone number during office hours:

Telephone number after office hours:

Information for Patients and Healthcare Professionals

Thalidomide can cause severe birth defects or death to an unborn baby therefore:

- Female patients of childbearing potential must always use effective contraception and Male patients with pregnant partners or partners of childbearing potential not using effective contraception must always use condoms (even if man has vasectomy)
- Female patients of childbearing potential must have pregnancy tests every 4 weeks, prior to each prescription, to ensure that they are not pregnant, except in the case of confirmed tubal sterilisation
- If a patient or partner of a patient suspects they are pregnant they must contact their **prescriber** immediately.

For complete information on the side effects of thalidomide, patients should read the Package leaflet and HCPs should read the summary of product characteristics

Information for Healthcare Professions:

Prescription details

Has the patient received counselling?:	Yes/No
Childbearing potential assessment:	WCBP / WNCBP / Male
If the patient is a WCBP is she using effective contraception?	Yes/No
If the patient is male, is he using condoms, if required?	Yes/No

A completed Prescription Authorisation Form must accompany each prescription to confirm that the patient continues to use effective contraception (if required) and, in the case of a WCBP, is having a pregnancy test every 4 weeks before each prescription to ensure they are not pregnant.

MHRA approval date 25/09/2020

BBBB2816